



---

## Q: What types of influenza vaccines are offered in Canada?

**A:** There are three types of influenza vaccines offered in Canada: inactivated influenza vaccines (IIV), the recombinant influenza vaccine (RIV), and live attenuated influenza vaccines (LAIV).

- **Inactivated influenza vaccines (IIV)** use an inactivated (killed) version of the influenza virus in the vaccine. They are given as an injection.
- **Recombinant influenza vaccine (RIV)** teaches your body to recognize a small protein on the surface of the influenza virus. It is given as an injection.
- **Live attenuated influenza vaccines (LAIV)** use an attenuated (weakened) form of the live influenza virus in the vaccine. They are given as a nasal spray.

IIV, RIV, and LAIV all teach your body how to recognize the influenza virus, so you will be able to fight it off in case you are infected.

---

## Q: Where can I receive an influenza vaccine?

**A:** Immunization programs are available in all provinces and territories. Contact your doctor, nurse, pharmacist, or local public health office about vaccination programs available in your area.

---

## Q: When should I be immunized?

**A:** The best time to get your influenza vaccine (flu shot) is in October, before influenza season starts (typically in November). You can get a flu shot anytime between October and March, but sooner is always better.

---

## Q: How long does it take for an influenza vaccine to work?

**A:** It takes about two weeks for an influenza vaccine to fully work. If you are exposed to the influenza virus within those two weeks, or shortly before your influenza vaccine (flu shot), you could catch influenza. However, this is not because of the vaccine.

---

## Q: Can I get influenza from the vaccine?

**A:** No. You cannot get influenza from an influenza vaccine.

---

## Q: How effective are the influenza vaccines?

**A:** The effectiveness of an influenza vaccine (its ability to prevent influenza illness) can range widely from season to season and from person to person. It can vary based on:

- a person's age (for example, persons 65+ may have a less robust immune response to the vaccine)
- a person's health (for example, persons with a weakened immune system may have a less robust immune response to the vaccine)
- how well the influenza vaccines match the strains of influenza circulating in a given influenza season

During years when there is a good match between the influenza vaccines and circulating viruses, it's more likely you will have good protection from severe influenza illness. In years where there is a mismatch, the vaccine may not be as effective, but it still provides some protection.

---

## Q: How safe are influenza vaccines?

**A:** Influenza vaccines have been safely used in Canada since the 1940s. They have been proven safe and can benefit people of all ages.

## **Q: Who should be immunized against influenza?**

**A:** All Canadians 6 months of age and older.

In particular, people at high risk of influenza infection and influenza-related complications, including:

- all pregnant persons (the risk of influenza-related hospitalization increases with length of gestation, i.e., it is higher in the third trimester than in the second)
- all children 6 to 59 months of age
- adults 65+
- Indigenous peoples
- people living in nursing homes and other chronic-care facilities
- children and adults with chronic health conditions such as:
  - cardiac or lung disorders, including asthma
  - diabetes and other metabolic diseases
  - cancer
  - immune-compromising conditions
  - kidney disease
  - anemia or hemoglobinopathy
  - morbid obesity (body mass index [BMI] of 40 and over)
  - children 6 months to 18 years of age with chronic conditions treated for long periods with acetylsalicylic acid
  - neurological or neurodevelopment conditions
- health care providers who could transmit influenza to people at high risk of influenza complications
- people who provide essential community services

---

## **Q: Is there anyone who shouldn't get an influenza vaccine?**

**A:** Most people can safely receive the influenza vaccine, but it may not be recommended for some individuals.

## **Influenza vaccination is NOT recommended in the following cases:**

- if you have had an anaphylactic reaction to a previous dose, or to any of the vaccine components
  - If you have had an anaphylactic reaction to one of the components in a specific influenza vaccine, you may be eligible to receive a different influenza vaccine where the component you are allergic to is absent. This must be done in consultation with an allergy expert.
- if you have had an onset of Guillain-Barré syndrome within six weeks of getting an influenza vaccine.

## **As well, it is recommended that you do not receive live attenuated influenza vaccines (LAIV) in the following cases:**

- if you have an immunocompromising condition (weakened immune system) [except children with stable HIV infection on antiretroviral therapy (ART)]
- if you have severe asthma, due to increased risk of wheezing after administration
- if you have received medical assistance due to wheezing seven days or less before your influenza vaccine (flu shot) appointment (there is an increased risk of wheezing after receiving LAIV)
- if your child is younger than 24 months (due to increased risk of wheezing after receiving LAIV)
- if you are pregnant (please see [Can I get immunized against influenza if I am pregnant?](#))
- if you are on antiretroviral therapies that target the influenza virus (within certain timeframes)

Talk to your healthcare professional if you have questions about getting LAIV or other influenza vaccines.

---

## Q: Are there side effects from influenza vaccines?

**A:** Influenza vaccines are very safe, but as with any medication, side effects can occur. Most people who get an influenza vaccine have mild side effects (such as soreness, redness, or swelling in the area where the influenza vaccine was given) or no side effects at all. Some people may get a fever or muscle aches that start shortly after immunization, which may last one to two days. Allergic reactions that can be life-threatening are extremely rare. If they do occur, it is within a few minutes to a few hours following vaccination. Your doctor, nurse, or pharmacist will ask you to wait 15 minutes following your vaccination, to watch for possible reactions.

---

## Q: Can influenza vaccines be given at the same time as other vaccines?

**A:** Yes. Other vaccines can be received at the same time as an influenza vaccine. For example, you can receive your COVID-19 vaccine at the same time as an influenza vaccine. Talk with your doctor, nurse, pharmacist, or local public health office about receiving more than one vaccine at the same time.

---

## Q: Can I get immunized against influenza if I am pregnant?

**A:** Yes. Influenza vaccines have been given to pregnant persons for decades and have a good safety record. There is no evidence that influenza vaccines harm the pregnant parent or fetus.

However, live attenuated influenza vaccines (LAIV) are not recommended for those who have weakened immune systems. Pregnancy can cause changes in your body that can affect your immune response.

**Therefore, you should NOT receive LAIV if you are pregnant.**



---

## Q: Can I get immunized against influenza if I am breastfeeding?

**A:** Yes. Influenza vaccines are considered safe and are recommended for all breastfeeding individuals.

---

## Q: Are children at an increased risk of developing neurological disorders like autism or attention-deficit hyperactivity disorder (ADHD) if they receive an influenza vaccine?

**A:** No. Multiple studies have shown that there is no link between vaccines and autism or attention-deficit/hyperactivity disorder (ADHD).

---

## Q: Why do some influenza vaccines contain thimerosal if other vaccines do not contain it?

**A:** Thimerosal is added to influenza vaccines in multi-dose vials to keep them sterile. It acts as a preservative, since multi-dose vials are used more than once. Single-dose influenza vaccines are used only once and do not need a preservative. There is no evidence that the use of thimerosal in vaccines causes harm.

---

## Q: Where can I get more advice on influenza immunization?

**A:** Contact your local public health office, doctor, nurse, or pharmacist. You can also read more about influenza immunization on the following websites:

- **Immunize Canada:** <https://immunize.ca>
- **Public Health Agency of Canada:** <https://www.canada.ca/en/public-health/services/diseases/flu-influenza.html>